

Morphological Transformation of BALB/3T3 Mouse Embryo Cells
Conducted with Test Article Extracts

CONCLUSION

The negative and positive controls fulfilled the requirements for determination of a valid test.

Under the conditions of the assay described in this report, saline and DMSO extracts of BIOCELL® Saline-filled Mammary Implant, Style 168 did not induce a statistically significant level of morphological transformation in the absence or presence of S-9 activation. BIOCELL® Saline-filled Mammary Implant, Style 168, was concluded to be negative in the BALB/3T3 cell transformation assay system in the absence and presence of exogenous metabolic activation.

REFERENCES

- IARC/NCI/EPA Working Group. 1985. Cellular and Molecular Mechanisms of Cell Transformation and Standardization of Transformation Assays of Established Cell Lines for the Production of Carcinogenic Chemicals: Overview and Recommended Protocols. *Cancer Res.* 45:2395-2399.
- Heidelberger, C., Freeman, A.E., Pienta, R.J., Slavak, A., Berman, J.S., Castro, B.C., Dunkel, V.C., Francis, M.W., Kakunaga, T., Little, J.R., and Schechtman, L.M. 1983. Cell transformation by chemical agents - a review and analysis of the literature. A report of the U.S. Environmental Protection Agency Geno-Tox Program. *Mutation Res.* 114:283-385.
- Kakunaga, T. 1973. A quantitative system for assay of malignant transformation by chemical carcinogens using a clone derived from BALB/3T3. *Int. J. Cancer* 12:463-473.
- McCarvill, J.T., Luter, R.A., Schechtman, L.M., Kouri, R.E., and Putman, D.L. 1990. Morphological Transformation of BALB/3T3 Cells by Various Procarcinogens in the Presence of a Rat Liver S9 Activation System. *Environ. Mol. Mutagenesis* 16:304-310.